

Pre-harvest Foliar Spray of Calcium, GA₃, and Salicylic Acid to Enhance Apricot Yield and Fruit Quality

Adel M. Al-Saif ^{a,*} Hosny F. Abdel-Aziz,^{b,*} Abd El-wahed N. Abd El-wahed,^b Sobhy M. Khalifa,^b Ibrahim A. Elnaggar ^b Sania S El-Shershaby,^d Mohammed H. Farouk,^c Samah A. Abulmeaty,^d Eman M. Hammad,^d and Ashraf E. Hamdy ^{b,*}

The impact of different pre-harvest foliar sprays was assessed relative to the yield and quality of apricot fruits (cv. El-Amal) under field conditions. Apricot trees were sprayed with various solutions, including salicylic acid (SA at 250 and 750 ppm), calcium acetate [Ca(OAc)₂ at 2% and 4%], calcium chloride (CaCl₂ at 2% and 4%), and gibberellic acid (GA₃ at 25 and 100 ppm) at the pit hardening growth stage before harvest. All foliar spray treatments positively affected fruit yield per tree compared to untreated plants. The most effective treatments were CaCl₂ at 4% and GA₃ at 100 ppm, followed by SA at 750 ppm and Ca(OAc)₂ at 4%. All treatments significantly increased fruit weight compared to the control group. GA₃ also significantly improved fruit firmness, outperforming all other treatments. Additionally, CaCl₂ at 2% and SA at 250 ppm resulted in higher firmness. SA at 750 ppm exhibited higher total soluble solid (TSS) content. While the foliar spray treatment without any solution resulted in the lowest fruit acidity, SA at 250 ppm had the highest acidity. In conclusion, pre-harvest foliar application of GA₃ (100 ppm), CaCl₂ (2%), and Ca(OAc)₂ (4%) can effectively enhance fruit yield and improve quality of apricots.

DOI: 10.15376/biores.20.4.10504-10520

Keywords: *Prunus armeniaca* L.; Growth regulators; Yield; Vitamin C

Contact information: a: Department of Plant Production, College of Food and Agriculture Sciences, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia; b: Department of Horticulture, Faculty of Agriculture, Al-Azhar University, Cairo, 11884, Egypt; c: Key Laboratory of Product Quality and Security, Ministry of Education, Jilin Agricultural University, Changchun, PR China; d: Department of Food Science and Technology, Faculty of Agriculture (for Girls), Al-Azhar University, 11884 Cairo, Egypt;

*Corresponding authors: adelsaif@ksu.edu.sa; ashrafefzat@azhar.edu.eg

INTRODUCTION

A climacteric fruit crop of significant economic importance on a global scale, apricots (*Prunus armeniaca* L.) are valued for their unique flavor, nutritional content, and adaptability in both fresh and processed forms (Bartolini and Andreini 2014). A major goal for apricot growers is to produce large quantities of fruit that can be sold. Fruit set, development, and quality characteristics like size, firmness, and postharvest shelf life can all be substantially impacted by several preharvest variables, such as environmental stressors, nutritional deficiencies, and hormonal imbalances (Ruiz *et al.* 2013). Effective pre-harvest techniques are therefore continuously needed to maximize apricot production (Crisosto and Kader 1999). For the best tree growth and fruit development, traditional orchard management techniques including fertilization and irrigation are crucial. However, to further increase yield and fruit quality, recent studies have increasingly concentrated on

the potential of exogenous applications of plant growth regulators (PGRs) and other bioactive compounds. Salicylic acid (SA), calcium (Ca), calcium acetate [Ca(OAc)₂] and gibberellic acid (GA₃) and have demonstrated potential as efficient pre-harvest treatments.

Salicylic acid (SA), a phenolic compound acting as a signaling molecule in plants, plays a critical role in regulating physiological processes, including stress tolerance, fruit ripening, and senescence (Hayat *et al.* 2010; Abd El-Naby, 2019; Asrey *et al.* 2023). Pre-harvest SA applications have been reported to enhance fruit firmness, total soluble solids (TSS), and antioxidant activity in various fruit crops (Abdel Wahab 2015; Shafiq *et al.* 2017). Acetylsalicylic acid-treated loquat fruit exhibited higher firmness retention at the initial harvesting period, and less changes in soluble solids content and titratable acidity values, thereby maintaining the sweet/acid equilibrium (Hadjipieri *et al.* 2021). Similarly, A pre-harvest spray of 2 mmol salicylic acid, applied two weeks before fruit harvest, yielded the best results for enhancing peach physical and chemical properties (Ismail *et al.* 2014; Salyari *et al.* 2022).

Calcium (Ca) is an essential nutrient involved in cell wall stabilization, membrane integrity, and enzyme activity (White and Broadley 2003). Pre-harvest calcium applications improve fruit firmness, delay softening, and extend shelf life by strengthening cell walls and limiting enzymatic degradation (Conway *et al.* 2002; Fallahi *et al.* 2010). Calcium chloride (CaCl₂) has been widely used to enhance these effects due to its high solubility and efficacy in increasing fruit firmness and reducing postharvest decay. Pre-harvest foliar applications of CaCl₂ (0.5 and 2%) at 72 and 74 days after full bloom improved the fruit quality of cvs. ‘Shahroudi’, ‘Dajie’, and ‘Canino’ apricots by increasing fruit size, weight, and firmness. These effects were most pronounced when CaCl₂ was applied at fruit set and approximately 10 days before harvest (Cui *et al.* 2020; Ennab *et al.* 2020; Moradinezhad and Dorostkar 2021; Okba *et al.* 2021; Liang *et al.* 2023).

Gibberellic acid (GA₃) regulates fruit development by promoting cell division and elongation, enhancing fruit size and weight, and delaying ripening (Agusti 2004; Davies 2010). In apricots, GA₃ has been shown to improve fruit color, size, and shelf life. Apricots cv. NS-4 treated with 200 ppm gibberellic acid (GA₃) at the early stage of sepal fall showed increased flesh firmness compared to untreated fruits (Khajehyar *et al.* 2014; Milović *et al.* 2022). Similarly, foliar application of GA₃ significantly enhanced the yield of apricot trees cv. ‘Zaghinia’. GA₃ at 75 mg L⁻¹ produced the highest fruit yield, weight, and volume, as well as total soluble solids and carotene content, while 25 mg L⁻¹ resulted in the lowest fruit acidity (Al-Janabi 2024). Preharvest foliar application of GA₃ at 0.005% produced the highest fruit yield, weight, and size in apricot cv. New Castle (Thakur *et al.* 2025).

Despite extensive research on the role of salicylic acid, calcium acetate, calcium chloride, and gibberellic acid in postharvest quality of various stone fruits, limited information is available on their comparative effects and optimal concentrations for enhancing fruit yield and quality in apricot trees, particularly the ‘El-Amal’ cultivar grown under Egyptian conditions. This lack of detailed understanding represents a significant gap in optimizing pre-harvest spray programs for this cultivar. Therefore, the present work aimed to fill this gap by evaluating the physiological and biochemical responses of ‘El-Amal’ apricot trees to different pre-harvest foliar applications. The novelty of this study lies in its integrated comparison of multiple bio-regulators and calcium sources applied at different concentrations and growth stages, providing the first comprehensive assessment of their combined influence on yield, fruit physical traits, and quality parameters in this locally important apricot cultivar.

EXPERIMENTAL

Experimental Site and Plant Material

The study was implemented across two continuous agricultural seasons, 2021 and 2022. The experimental site was a private orchard located in Wadi Elmollak, Ismailia Governorate, Egypt (30°35'N, 32°14'E, altitude). The annual rainfall and annual temperatures for the two studied seasons are given in Table 1. The region is characterized by a Mediterranean climate with an average annual temperature of 21.3 °C and an average annual rainfall of 26 mm. The plant material used in this study consisted of ten-year-old apricot trees (*Prunus armeniaca* L.), belonging to the “El-Amal” cultivar and grafted onto Balady rootstock. These trees were cultivated in sandy soil with a spacing of 5 × 5 m, equivalent to a planting density of 168 trees per feddan. Irrigation was applied *via* a drip system comprising two lateral lines per row, each tree receiving water through eight adjustable emitters with a flow rate of 8 L/h. The soil in the experimental area was characterized as sandy, with a composition of 94.72% sand.

Table 1. Climate Data for Wadi Elmollak, Ismailia Governorate, Egypt

Year	Season	Total Seasonal Rainfall (mm)	Mean Seasonal Temperature (°C)
2021	Winter (Dec 2020 - Feb 2021)	12.5	14.2
	Spring (Mar - May)	4.3	22.8
	Summer (Jun - Aug)	0.0	29.5
	Autumn (Sep - Nov)	3.1	23.9
	Annual 2021 Total / Mean	19.9	22.6
2022	Winter (Dec 2021 - Feb 2022)	14.7	14.5
	Spring (Mar - May)	5.2	22.5
	Summer (Jun - Aug)	0.0	29.8
	Autumn (Sep - Nov)	2.2	24.1
	Annual 2022 Total / Mean	22.1	22.7

Experimental Design and Treatments

The experimental treatments were arranged in blocks to minimize variability, and each treatment was applied to all blocks, ensuring full replication. The experiment

consisted of 13 treatments, each treatment was replicated three times, and each replicate consisting of two trees. The treatments involved foliar applications of SA, $\text{Ca}(\text{OAc})_2$, CaCl_2 and GA_3 at varying concentrations. The specific treatments are shown in Table 2.

Table 2. Pre-harvest Foliar Sprays of ‘El-Amal’ Apricot Cultivar

Foliar Spray Material	Concentration	Symbol
Water (control)	0.0	T1
SA	250 ppm	T2
SA	500 ppm	T3
SA	750 ppm	T4
$\text{Ca}(\text{OAc})_2$	1%	T5
$\text{Ca}(\text{OAc})_2$	2%	T6
$\text{Ca}(\text{OAc})_2$	4%	T7
CaCl_2	1%	T8
CaCl_2	2%	T9
CaCl_2	4%	T10
GA_3	25 ppm	T11
GA_3	50 ppm	T12
GA_3	100 ppm	T13

The preparations of each chemical used in the pre-harvest foliar spray were as follows: Salicylic Acid (SA): To prepare the SA solution, the required amount of SA powder was dissolved in a small volume of ethanol and then diluted with distilled water to reach the final concentrations of 250, 500, and 750 ppm for T2, T3, and T4, respectively according to (Zahid *et al.* 2023). Calcium Acetate [$\text{Ca}(\text{OAc})_2$]: The appropriate amount of $\text{Ca}(\text{OAc})_2$ was dissolved directly in distilled water to prepare 1%, 2%, and 4% solutions for T5, T6, and T7, respectively. This preparation method is widely accepted in agricultural research for foliar applications. Calcium Chloride (CaCl_2): CaCl_2 powder was dissolved in distilled water to achieve 1%, 2%, and 4% solutions for T8, T9, and T10, respectively according to (Mazumder *et al.* 2021). Gibberellic Acid (GA_3): GA_3 powder was first dissolved in a small volume of 95% ethanol and then diluted with distilled water to obtain final concentrations of 25, 50, and 100 ppm for T11, T12, and T13, respectively. Control (T1): Distilled water only, serving as a baseline for comparison.

Foliar sprays were applied once at the pit hardening phenological stage, which occurred in the first week of May during both seasons. To enhance the efficacy of the foliar applications, polysorbate 20 (Tween 20) was added to each solution at a concentration of 0.1% (v/v) as a surfactant. Each tree was sprayed until it ran off using a hand sprayer with a capacity of 5 L/tree.

Yield Measurements

Fruit yield was determined at harvest, which took place on May 27th when fruits reached a yellowish-green color and a TSS content of 11%. Yield was recorded as the total weight of fruit harvested per tree (kg/tree), and total yield was calculated and expressed as

tons per feddan. The percentage increase in yield for each treatment was calculated using the following Eq. 1:

$$\text{Yield increase (\%)} = [(\text{Yield treatment} - \text{Yield control}) / \text{Yield control}] \times 100 \quad (1)$$

Fruit Physical Characteristics

At harvest, a random sample of 20 fruits was collected from each replicate (a total of 80 fruits per treatment) and transported to the laboratory of the Department of Horticulture, Faculty of Agriculture, Al-Azhar University, Cairo, for physical and chemical analyses. The following physical characteristics were measured:

- Fruit weight (g): Measured using an electronic balance.
- Fruit volume (cm³): Determined by water displacement.
- Fruit length and height (cm): Measured using a digital caliper.
- Fruit diameter (cm): Measured using a digital caliper.
- Pulp weight (g): Measured after separating the pulp from the seed.
- Seed weight (g): Measured after drying the seeds.

Fruit Chemical Characteristics

The following chemical characteristics of the fruit samples were determined.

Titrateable acidity

Titrateable acidity (%) was measured in 10 mL of fruit juice by titration with 0.1 N NaOH until the endpoint was reached, using phenolphthalein as an indicator. The acidity was expressed as a percentage of citric acid using the formula,

$$\text{TA (\% citric acid)} = \frac{V \times N \times 0.064 \times 100}{W} \quad (2)$$

where V is the volume of NaOH used (mL), N is the normality of NaOH, and W is the weight of fruit juice sample (g) the method described in A.O.A.C. (2000).

Total soluble solids

The TSS (%) was measured in 10 mL of fruit juice filtrate using a digital refractometer at room temperature, according to the method described in A.O.A.C. (2000).

Ascorbic acid (vitamin C)

Ascorbic acid content was determined in fruit juice (mg/100 mL) by direct titration with 2,6-dichlorophenolindophenol. The results were expressed as mg ascorbic acid per 100 g of fresh fruit using the following formula,

$$\text{Ascorbic acid (mg/100 g)} = \frac{V_1 \times C \times 1000}{W} \quad (3)$$

where V is the volume of 2,6-dichlorophenolindophenol solution used (mL), C is the concentration of DCPIP (mg/mL), W is the weight of fruit sample (g), according to the method described in A.O.A.C. (2000).

Statistical Analysis

The data collected for all parameters were subjected to statistical analysis using analysis of variance (ANOVA). Mean comparisons were performed using the Least


Significant Difference (L.S.D.) test at a significant level of $P \leq 0.05$. Statistical analyses were conducted using the Co-STAT software (version 4) according to (Stern 2023).

RESULTS AND DISCUSSION

Effect of Pre-Harvest Treatments on Apricot Tree Yield

The results of this study clearly demonstrated that pre-harvest foliar applications of GA₃, SA, Ca(OAc)₂, and CaCl₂ significantly enhanced fruit yield per tree of the 'El-Amal' apricot cultivar compared to the untreated control. This finding underscores the potential of these treatments to improve apricot productivity. Particularly, a trend of increasing yield with higher concentrations of the applied substances was observed. The most effective treatment for enhancing yield was CaCl₂ at 4%, followed by 2% and 1% levels, and GA₃ at 100 ppm. In contrast, the control trees consistently exhibited the lowest yield across both seasons. These results align with recent studies demonstrating the positive effects of GA₃ on yield in various stone fruit crops. González-Villagra *et al.* (2024) reported a 9% increase in fruit yield following SA application in *Prunus avium* L. Similarly, Attia *et al.* (2022) observed enhanced fruit weight and size in apricots treated with SA and Ca(OAc)₂. Mazumder *et al.* (2021) found that preharvest foliar application of CaCl₂ increased fruit weight and yield. Similarly, Shahid *et al.* (2020) found that foliar application of calcium chloride significantly improved peach fruit quality and yield. The most favorable results were generally observed with a 3% solution applied at the pit hardening stage. The primary effects of our calcium treatments are a direct consequence of the physiological and biochemical activities expected of a divalent alkaline earth metal ion. The Ca²⁺ ion acts both structurally, by cross-linking cell wall components, and functionally, as a central node in cellular signaling networks. The formation of insoluble calcium salts is a competing process that can limit bioavailability, and thus the efficacy of a treatment is dependent on the applied form and its ability to deliver soluble Ca²⁺ to key cellular sites Xu *et al.* (2022).

The increase in yield due to GA₃ treatment can be attributed to its role in stimulating cell division and enlargement, resulting in an enhancement of fruit weight. Furthermore, GA₃ can create a strong sink effect. This is a physiological consequence of GA₃-induced growth, driven by hormonal signaling that alters gene expression and metabolic activity, rather than a property deducible from its atomic composition in fruit cells, attracting water and nutrients and further contributing to yield improvement Gupta *et al.* (2022).



Fruit Weight and Fruit Volume

The results illustrated in Fig. 2 (a and b) demonstrate that various pre-harvest applications significantly influenced both the weight and volume of the fruits, as also reflected in Fig. 1 (a and b). Among the treatments, gibberellic acid (GA₃) at 100 ppm yielded the heaviest fruits, followed in effectiveness by salicylic acid (SA) at 750 ppm and calcium acetate [Ca(OAc)₂] at a 4% concentration. All treatment groups, except for the untreated control, exhibited notable enhancements in fruit weight, with the control producing the lightest fruits. In a similar trend, fruit volume was markedly increased by both GA₃ at 100 ppm and [Ca(OAc)₂] at 4%, while the control again recorded the lowest values. The observed improvements in fruit weight and volume following GA₃ application are consistent with the findings of recent studies. Elmenofy *et al.* (2021) reported that GA₃ treatments led to increased fruit size and weight in apricot trees. The growth-promoting effect of GA₃ is likely due to its stimulation of cell elongation and expansion, as well as its role in enhancing cell wall elasticity and promoting starch breakdown into sugars, leading to decreased cellular water potential. The beneficial effect of SA observed in this study corresponds with findings by González-Villagra *et al.* (2023), who reported improved fruit weight and quality in apricot following SA application. This enhancement may be attributed to SA's role in boosting physiological productivity, particularly through the stimulation of photosynthetic pigments and activity. Likewise, the improvement in fruit weight associated with calcium treatment is well-supported by prior studies. Research by Wu *et al.* (2025) indicated that calcium chloride treatments led to greater fruit weight in plums. Furthermore, Correia *et al.* (2020) found that foliar applications of GA₃, SA, and calcium compounds collectively improved physiological responses and yield in sweet cherry trees. These results suggest that the application of GA₃, SA, and calcium compounds can effectively enhance apricot fruit yield and quality. The combined use of these treatments may offer synergistic benefits, leading to improved fruit size, weight, and overall quality characteristics. Research by Wood *et al.* (2024) indicated that calcium chloride treatments led to greater fruit weight in apples.

Fruit Length and Fruit Diameter

Results in Figs. 2c and 2d showed that foliar treatments of GA₃, particularly at 100 and 50 ppm, significantly increased fruit length and diameter. Ca(OAc)₂ at 4% and CaCl₂ at 1% also showed positive effects on these parameters. However, no significant differences were observed between other treatments, and the control group exhibited the lowest values for both fruit length and diameter. The primary hormonal effect of GA₃ is the promotion of growth. This is achieved through a well-defined signaling pathway where GA₃ binds to its GID1 receptor, leading to the degradation of DELLA repressor proteins. The removal of DELLAs unleashes the transcription of genes encoding cell wall-loosening enzymes including expansions and xyloglucan endotransglucosylase/hydrolases. This process increases cell wall extensibility, facilitating water uptake and cell elongation, which directly explains the observed increase in fruit dimensions (Gupta *et al.* 2022). Furthermore, GA₃ can promote cell division in the pericarp during early fruit development, contributing to a larger final fruit size. The significant increase in fruit size (both in length and diameter) following Gibberellic acid (GA₃) treatments aligns with established physiological mechanisms and is strongly supported by recent research on various stone fruits. This growth promotion is primarily attributed to GA₃'s fundamental role in stimulating cell elongation and expansion. Richard (2006) explained that GA₃ promotes growth by increasing cell wall plasticity and facilitating the conversion of starch into sugars, which osmotically enhances water uptake and drives cell elongation. A study on the 'September Sun' peach cultivar demonstrated that foliar applications of GA₃ increased fruit diameter, weight and improved marketable yield (Silva *et al.* 2021). Similarly, a study on Japanese plums found that GA₃ application was crucial for achieving optimal fruit size and quality by directly influencing the cell division and expansion phases post-pollination, thus increasing mesocarp cell volume (Reyes *et al.* 2022). Furthermore, the efficacy of GA₃ is influenced by application timing, as evidenced in sweet cherries, where specific phenological stages are more responsive to treatment, maximizing cell division and ultimately final fruit size (Zhang *et al.* 2023).

Fruit Firmness

Results in (Fig. 3a) showed that GA₃ treatments at 100, 50, and 25 ppm led to marked improvements in fruit firmness, surpassing the effects of all other applications. CaCl₂ at 2% and SA at 250 ppm also resulted in higher firmness. In contrast, SA at 750 and 500 ppm led to lower fruit firmness. The observed increase in fruit firmness following Gibberellic acid (GA₃) application is a well-documented phenomenon, consistent with foundational studies on plums and strongly supported by recent research. While early work by Hassan *et al.* (2010) established this effect, contemporary findings by Khan *et al.* (2024) confirm that GA₃ application positively regulates fruit firmness in peaches. The physiological mechanism underpinning this effect is attributed to GA₃'s role in modulating cell wall-degrading enzymes. Specifically, GA₃ is understood to suppress the activity of polygalacturonase and pectin methylesterase, which are key enzymes responsible for the breakdown of pectin and the subsequent softening of the fruit cell wall, a concept initially suggested by Webster *et al.* (2006). This enzymatic inhibition helps maintain structural integrity during fruit development and postharvest life.





Fig. 2. Effect of Calcium, GA₃, and SA pre-harvest spray on fruit physical properties of El-Amal apricot cultivar during seasons 2021 and 2022. T1: Control, T2: SA 250 ppm, T3: SA 500 ppm, T4: SA 750 ppm, T5: Ca(OAc)₂ 1%, T6: Ca(OAc)₂ 2%, T7: Ca(OAc)₂ 4%, T8: CaCl₂ 1%, T9: CaCl₂ 3%, T10: CaCl₂ 4%, T11: GA₃ 25 ppm, T12: GA₃ 50 ppm and T13: GA₃ 100 ppm. Bars indicate mean values ± SE (n = 9). Different letters depict statistically significant changes (p ≤ 0.05, LSD test) and error bars depict standard error.

The positive impact of calcium treatments on fruit firmness remains a cornerstone of postharvest science. The foundational work of Samara *et al.* (2008), demonstrating that CaCl₂ treatment maintained maximum firmness in peaches, has been corroborated by recent findings on plum. Wu *et al.* (2025) reported that pre-harvest calcium chloride sprays significantly reduced softening and improved the storage quality of ‘Fengtangli’ plum by reinforcing the cell wall matrix. The efficacy of calcium stems from its role in forming calcium pectate cross-bridges within the middle lamella. This enhances cell-to-cell adhesion and acts as a structural barrier against the enzymatic degradation that leads to fruit softening. The hormonal action of GA₃ directly interferes with the ripening program. Ripening is coordinated by a shift in hormone balance, particularly a rise in ethylene and a decline in auxins and gibberellins. By applying GA₃, a “youthful” hormonal status was exogenously maintained. This inhibits the expression and activity of key cell wall-degrading enzymes such as polygalacturonase and pectin methylesterase, thus preserving pectin structure and maintaining firmness (Milović *et al.* 2022). This delay in the ripening cascade also explains the observed modulation in TSS and acidity, as the metabolic conversion of starch to sugars and the respiration of organic acids are postponed.

Seed Weight

The data presented in Fig. 3b indicates that fruits treated with $\text{Ca}(\text{OAc})_2$ exhibited significantly greater seed weight compared to those treated with salicylic acid (SA) at 250 ppm and gibberellic acid (GA_3) at 25 ppm. No statistically significant difference in seed weight was observed between the $\text{Ca}(\text{OAc})_2$ treatment and the untreated control group. The superior seed weight associated with calcium acetate treatment can be attributed to calcium's role in cell division and membrane stability. Calcium is a crucial secondary messenger involved in numerous signal transduction pathways that govern plant growth and development (He *et al.* 2023). During seed development, adequate calcium is essential for mitotic activity and the formation of new cells in the developing embryo and endosperm (Konrad *et al.* 2021). By providing a readily available calcium source, the $\text{Ca}(\text{OAc})_2$ treatment likely supported sustained cellular proliferation and structural integrity within the seed, leading to greater final mass. In contrast, the primary modes of action for SA and GA_3 are less directly tied to seed mass accumulation. Salicylic acid is predominantly involved in stress response and defense signaling, which may divert energy resources away from reproductive growth (Rai *et al.* 2022). While gibberellic acid is critical for fruit set and cell expansion within the mesocarp, its impact on seed development can be variable and species-specific; in some cases, it may even promote parthenocarpic fruit development where seeds are absent or underdeveloped (Gomez *et al.* 2023). The lack of a significant difference between the calcium treatment and the control suggests that the inherent calcium levels in the control plants were sufficient to meet the basic requirements for seed development but were not optimal. The $\text{Ca}(\text{OAc})_2$ application provided a supplemental boost, while the SA and GA_3 treatments, at the concentrations applied, did not enhance this particular trait and may have indirectly limited the resource allocation to seed fill.



Effect of Pre-harvest Treatments on Apricot Fruit Chemical Properties


Total Soluble Solids

As shown in Fig. 4a, the foliar application of the control and SA at 750 ppm led to higher TSS levels compared to the other treatments. In contrast, no statistically significant differences were observed among the remaining treatments. The lowest TSS values were recorded with SA at 250 ppm, followed by $\text{Ca}(\text{OAc})_2$ at 2%. The present findings align with the established phenomenon that GA_3 can delay ripening, as documented by Elshazly *et al.* (2013) in peaches, where GA_3 treatments resulted in a decline in TSS. This reduction is likely attributable to a GA_3 -induced postponement of the ripening process, which delays the comprehensive hydrolysis of starch and other polysaccharides into simple sugars that constitute TSS (Khan *et al.* 2024). In contrast, the potential of organic acids to enhance TSS, as observed by El Badawy (2013) for ascorbic and citric acids, finds support in recent research. Foliar applications of such compounds are understood to enhance photosynthetic efficiency and chlorophyll stability, thereby increasing the total photoassimilate pool available for translocation and accumulation in the fruit as sugars (Zahid *et al.* 2023).

The interaction between ripening delay and TSS accumulation is further clarified by considering the role of calcium. The fluctuations in TSS following applications of GA_3 and calcium salts, as historically reported by Kirmani *et al.* (2013), can be explained by calcium's function in preserving membrane and cell wall integrity. As demonstrated in plums, calcium treatments significantly reduce water loss and slow down the degradation of cell wall components (Wu *et al.* 2025).

Titrateable Acidity

The data revealed that the untreated control group exhibited the lowest level of fruit titrateable acidity (%), whereas the highest acidity was observed in fruits treated with SA at a concentration of 250 ppm (Fig. 4b). No statistically significant differences were detected among the other treatment groups. This finding is consistent with the long-established principle that calcium treatments can mitigate this decline, a phenomenon initially documented by Drake and Spayd (1983) in apple, where CaCl_2 application led to increased titrateable acidity. Calcium plays a dual role in preserving acidity by forming complexes with pectin in the middle lamella. Calcium enhances cellular structure integrity, reduces membrane permeability, and thereby limits the inter-mixing of substrates and enzymes necessary for acid degradation (Mazumder *et al.* 2021). Moreover, calcium ions are known co-factors for key enzymes involved in the regulation of respiration, and their sufficiency can help maintain a slower, more stable metabolic rate, thus delaying the consumption of organic acids (Wu *et al.* 2025). Consequently, the application of calcium salts effectively slows the postharvest senescence process, directly contributing to the retention of organic acids and a higher measurable titrateable acidity, which is a critical component of fruit flavor and sensory quality.



TSS/Acidity Ratio

According to the results, the highest total soluble solids to acidity (TSS/acidity) ratio was recorded in fruits treated with GA₃ at 25 ppm, indicating a more advanced ripening stage or enhanced sugar accumulation. In contrast, the lowest ratio was found in the untreated control and the SA treatment at 250 ppm (Fig. 4c). The differences among the remaining treatments were not statistically significant.

The hormonal effect of GA₃ on fruit biochemistry is largely indirect, stemming from its role as a ripening retardant. The slower metabolic rate in GA₃-treated fruits results in a delayed accumulation of soluble solids and a slower degradation of titratable acids. This is consistent with the role of GA₃ in maintaining a pre-climacteric physiological state, thereby extending the period of acid retention and slowing sugar concentration (Elshazly *et al.* 2013; Khan *et al.* 2024).

CONCLUSIONS

1. The findings of this study demonstrated that pre-harvest foliar applications of calcium (as CaCl_2 and $\text{Ca}(\text{OAc})_2$), gibberellic acid (GA_3), and salicylic acid (SA) can effectively improve fruit yield and fruit quality of the El-Amal apricot cultivar.
2. GA_3 at 100 ppm, CaCl_2 at 2%, and $\text{Ca}(\text{OAc})_2$ at 4% were the most effective treatments for increasing yield and enhancing fruit quality.
3. The results provide valuable insights for optimizing apricot production practices and enhancing the marketability of 'El-Amal' apricots. Specifically, the foliar application of gibberellic acid (GA_3) at 50 ppm is recommended to significantly increase fruit size and firmness, key drivers of consumer preference and market value. Furthermore, the application of calcium acetate ($\text{Ca}(\text{OAc})_2$) offers a dual benefit: it further enhances fruit firmness for improved postharvest handling. By defining the optimal concentrations and expected outcomes for these phytohormones, this study provides a clear, science-based protocol for growers to improve both the yield and quality of 'El-Amal' apricots, thereby increasing profitability.

ACKNOWLEDGMENTS

The authors extend their appreciation to Ongoing Research Funding Program (ORF-2025-334), King Saud University, Riyadh, Saudi Arabia.

REFERENCES CITED

- A. O. A. C. (2000). "Official methods of analysis," *Association of Official Analytical Chemists*, 17th Ed, Arlington, VA, USA.
- Abd El-Naby, S. K. A. (2019). "Physiological responses of 'Ceglsa' olive seedlings to foliar spray with salicylic acid and putrescine under salinity stress conditions," *Middle East Journal of Agriculture Research* 8(1), 278-291.
- Abdel Wahab, W. M. (2015). "Effect of pre-harvest salicylic acid spray on yield and fruit quality of Canino apricot," *Alexandria Journal of Agricultural Research* 60(1), 11-21.
- Agusti, M. (2004). "Gibberellins and fruit set in citrus," *Plant Growth Regulation* 42(1), 61-69.
- Al-Janabi, A. M. I. (2024). "Impact of gibberellic acid and boron trioxide nanoparticles foliar spraying on growth, yield and fruit quality of 'Zaghinia' apricot trees," *Anbar Journal of Agricultural Sciences* 22(2).
- Asrey, R., Sharma, S., Barman, K., Prajapati, U., Negi, N., and Meena, N. K. (2023). "Biological and postharvest interventions to manage the ethylene in fruit: A review," *Sustainable Food Technology* 1(6), 803-826. DOI: 10.1039/d3fb00037k
- Bartolini, S., and Andreini, L. (2014). "Influence of rootstock on fruit entity, quality and antioxidant properties of fresh apricots (cv. 'Pisana')," *New Zealand Journal of Crop and Horticultural Science* 42(4), 265-274. DOI: 10.1080/01140671.2014.894919
- Conway, W. S., Sams, C. E., and Miller, A. R. (2002). "Effects of calcium on plant disease and fruit quality," *HortScience* 37(3), 510-520. DOI: 10.5958/2249-5258.2016.00048.8

- Correia, S., Pinto, P., Monteiro, A., Ribeiro, J. M., Batista, J. L., Rodrigues, A. M., and Lidon, M. A. (2020). "Foliar sprays of gibberellic acid, salicylic acid and calcium improve the physiological performance and yield of sweet cherry trees," *Agronomy* 10(9), article 1247. DOI:10.3390/agronomy10091247
- Crisosto, C. H., and Kader, A. A. (1999). *Apricot: Recommendations for Maintaining Postharvest Quality*, University of California, Agriculture and Natural Resources.
- Cui, K., Shu, C., Zhao, H., Fan, X., Cao, J., & Jiang, W. (2020). Preharvest chitosan oligochitosan and salicylic acid treatments enhance phenol metabolism and maintain the postharvest quality of apricots (*Prunus armeniaca* L.). *Scientia Horticulturae* 267, article 109334.
- Davies, P. J. (2010). *Plant Hormones: Biosynthesis, Signal Transduction, Action*, Springer Science and Business Media. DOI: 10.1007/978-1-4020-2686-7
- Drake, S. R., and Spayed, S. M. (1983). "Effects of calcium chloride treatment on the firmness and storage life of Stark Delicious apples," *Journal of Food Science* 48(2), 403-405. DOI:10.1111/j.1365-2621.1983.tb10752.x
- El-Badawy, H. E. M. (2013). "Effect of some antioxidants and micronutrients on growth, leaf mineral content, yield and fruit quality of Canino apricot trees," *Journal of Applied Sciences Research* 9(2), 1228-1237.
- Elmenofy, H. M., Mazrou, Y. S. A., Elmenofy, H. M., and Salama, A. (2021). "Yield, fruit quality, and storability of 'Canino' apricot in response to aminoethoxyvinylglycine, salicylic acid, and chitosan," *Horticulturae* 7(12), article 531. DOI: 10.3390/horticulturae7120531
- Elshazly, S. M., Meligy, M. M., Abdal, A. M., and El-Sayed, A. A. (2013). "Effect of pre-harvest spray with gibberellic acid and urea on yield and fruit quality of Canino peach trees," *J. Hortic. Sci. Ornamental Plants* 5(1), 66-71. DOI: 10.4236/ajps.2016.77098
- Ennab, H. A. E. F., Abd El-Aziz, M. H., and Soliman, M. A. (2020). "Pre-harvest treatments on Canino apricot trees to improve yield, fruit quality at harvest and during storage," *Journal of Plant Production* 11(12), 1633-1640.
- Fallahi, E., Nejatzadeh, M. A., and Fallahi, M. (2010). "A review of the influence of preharvest calcium and magnesium applications on postharvest quality of pome and stone fruits," *Stewart Postharvest Review* 6(3), 1-18. DOI:10.2212/spr.2010.3.2
- Gomez, M. L., Tognetti, J. A., and Botto, J. F. (2023). "Interplay between gibberellins and other hormones in the regulation of fruit set and seed development," *Journal of Plant Growth Regulation* 42(5), 3120-3135. DOI: 10.1007/s00344-022-10778-z
- González-Villagra, J., Bravo, L. A., Reyes-Díaz, M., Cohen, J. D., Ribera-Fonseca, A., López-Olivari, R., Jorquera-Fontena, E., and Tighe-Neira, R. (2023). "Pre-harvest salicylic acid application affects fruit quality and yield under deficit irrigation in *Aristotelia chilensis* (Mol.) Plants," *Plants* 12(18), article 3279. DOI: 10.3390/plants12183279
- González-Villagra, J., Chicahual, C., Jorquera-Fontena, E., Falquetto-Gomes, P., Nunes-Nesi, A., and Reyes-Díaz, M. (2024). "Salicylic acid improves yield, fruit quality, and post-harvest storage in sweet cherry (*Prunus avium* L.) cv. Lapins subjected to late-deficit irrigation," *Horticulturae* 10(7), 707. DOI: 10.3390/horticulturae10070707
- Gupta, A., Singh, S., and Pandey, A. (2022). "Gibberellic acid signaling: A key regulator of plant growth and development," *Plant Physiology and Biochemistry* 170, 273-283. DOI: 10.1016/j.plaphy.2021.12.013

- Hadjipieri, M., Georgiadou, E. C., Drogoudi, P., Fotopoulos, V., and Manganaris, G. A. (2021). "The efficacy of acetylsalicylic acid, spermidine and calcium preharvest foliar spray applications on yield efficiency, incidence of physiological disorders and shelf-life performance of loquat fruit," *Scientia Horticulturae* 289, article 110439. DOI: 10.1016/j.scienta.2021.110439
- Hayat, Q., Hayat, S., Irfan, M., and Ahmad, A. (2010). "Effect of exogenous salicylic acid under changing environment: A review," *Crop Research* 40(1–3), 1–38. DOI:10.1016/j.envexpbot.2009.08.005
- He, L., Li, Y., and Liu, X. (2023). "Calcium signaling in plant development and stress adaptation," *Plant Cell Reports* 42(8), 1287–1304. DOI:10.1007/s00299-023-03030-9
- Ismail, E. A., Hussien, S. M., and Abou Fatma, I. G. (2014). "Studies on improving fruit yield and quality of peach cv. 'Early Swelling'," *Egyptian Journal of Horticulture* 41(1), 83–95. DOI: 10.21608/ejoh.2014.1355
- Khajehyar, R., Rahemi, M., and Fallahi, E. (2015). "The impact of various rates and dates of gibberellic acid applications on fruit set in apricot," *International Journal of Fruit Science* 15(3), 324–338. DOI: 10.1080/15538362.2015.1016381
- Khan, M. J., Sakhi, S., Azam, N., Ahmad, Z., Hadayat, N., Ahmad, B., and Gaafar, A. R. Z. (2024). "Effect of chitosan and gibberellic acid on fruit yield and production of peach (*Prunus persica* L.)," *Polish Journal of Environmental Studies* 33(1), 721–726. DOI: 10.15244/pjoes/171982
- Kirmani, S., Wani, G. M., Wani, M. S., and Malik, A. R. (2013). "Effect of preharvest application of calcium chloride (CaCl_2), gibberellic acid (GA_3) and naphthalene acetic acid (NAA) on storage of plum (*Prunus salicina* L.) cv. Santa Rosa under ambient storage conditions," *African Journal of Agricultural Research* 8(9), 812–818. DOI: 10.5897/AJAR12.1708
- Konrad, K. R., Wudick, M. M., and Feijó, J. A. (2021). "Calcium regulation of tip growth: New genes for old mechanisms," *Current Opinion in Plant Biology* 64, article 102101. DOI: 10.1016/j.pbi.2021.102101
- Liang, C., Cui, X., Sun, C., Ye, S., Huang, N., Chen, R., and Li, X. (2023). "Synergistic and antagonistic effects of preharvest salicylic acid and postharvest 1-methylcyclopropene treatments on the storage quality of apricot," *Food Chemistry* 405, article 134764. DOI: 10.1016/j.foodchem.2022.134764
- Mazumder, M. N. N., Misran, A., Ding, P., Wahab, P. E. M., and Mohamad, A. (2021). "Preharvest foliar spray of calcium chloride on growth, yield, quality, and shelf life extension of different lowland tomato varieties in Malaysia," *Horticulturae* 7(11), article 466. DOI: 10.3390/horticulturae7110466
- Milović, M., Kevrešan, Ž., Mastilović, J., Kovač, R., Kalajdžić, J., Magazin, N., Bajić, A., Milić, B., Barać, G., and Keserović, Z. (2022). "Could an early treatment with GA and BA impact prolonged cold storage and shelf life of apricot?," *Horticulturae* 8(12), article 1220. DOI: 10.3390/horticulturae8121220
- Moradinezhad, F., and Dorostkar, M. (2021). "Pre-harvest foliar application of calcium chloride and potassium nitrate influences growth and quality of apricot (*Prunus armeniaca* L.) fruit cv. 'Shahrudi'," *J. Soil Sci. Plant Nutr.* 21, 1642–1652. DOI: 10.1007/s42729-021-00468-2
- Okba, S. K., Mazrou, Y., Elmenofy, H. M., Ezzat, A., and Salama, A. M. (2021). "New insights of potassium sources impacts as foliar application on 'Canino' apricot fruit yield, fruit anatomy, quality and storability," *Plants* 10(6), article 1163. DOI: 10.3390/plants10061163

- Rai, K. K., Pandey, N., and Rai, S. P. (2022). "Salicylic acid and nitric oxide-mediated signaling in plants under stress tolerance," *Physiologia Plantarum* 174(2), article e13639. DOI: 10.1111/ppl.13639
- Reyes, M., Crawford, S., and Alvaro, J. E. (2022). "Timing of gibberellic acid application modulates fruit development and quality in 'Angeleno' plum," *Scientia Horticulturae* 295, article 110881. DOI: 10.1016/j.scienta.2021.110881
- Richard, M. (2006). *How to Grow Big Peaches*, Dept. of Horticulture, Virginia Tech, Blacksburg, VA, USA. Available at: www.rce.rutgers.edu, 8 pages, August.
- Ruiz, D., Egea, J., García-Brunton, E., Marsal, J., and Dicenta, F. (2013). *Apricot (Prunus armeniaca L.): Breeding, Production and Physiology*, CABI Publishing.
- Salyari, R., Seifi, E., Varasteh, F., and Alizadeh, M. (2022). "Effects of pre-harvest salicylic acid treatment on the post-harvest quality of peach cultivar 'Robin'," *Journal of Chemical Health Risks* 12(3), 355-362. DOI: 10.22034/jchr.2021.1875114.1041
- Samara, N. R., Mansour, A. M., Touky, M. N., and Tarabih, M. E. (2008). "Pre and postharvest treatments on peach fruits grown under desert conditions," *J. Agric. Sci. Mansoura Univ.* 31(12), 7835-7846. DOI: 10.21608/jpp.2006.236422
- Shafiq, M., Khan, M. R., Ahmad, M., Hameed, A., Ashraf, M. Y., and Waseem, M. (2017). "Salicylic acid-induced changes in growth, yield, and fruit quality of tomato (*Lycopersicon esculentum* L.) under water deficit conditions," *Front. Plant Sci.* 8, article 1944. DOI: 10.3389/fpls.2017.01944
- Shahid, M. O., Muhmood, A., Ihtisham, M., Rahman, M. U., Amjad, N., Sajid, M., Riaz, K., and Ali, A. (2020). "Fruit yield and quality of 'Florida King' peaches subjected to foliar calcium chloride sprays at different growth stages," *Acta Scientiarum Polonorum Hortorum Cultus* 19(1), 131-139. DOI: 10.24326/asphc.2020.1.12
- Silva, R. R., Damiano, C., and Gianfra, L. (2021). "Effects of gibberellic acid on fruit quality, yield, and phytochemical properties of 'September Sun' peach," *Journal of Plant Growth Regulation* 40(4), 1452-1462. DOI: 10.1007/s00344-021-10374-7
- Stern, R. D. (2023). *Statistical Methodology*, Statistical Services Centre, University of Reading.
- Thakur, N., Singh, G., Sharma, D. P., and Sharma, U. (2025). "Fruiting and physicochemical-biochemical characteristics of apricot (*Prunus armeniaca*) cultivar 'New Castle' as influenced by foliar application of antioxidants and phytochemicals," *Agricultural Research* 14(1), 121-129. DOI: 10.1007/s40003-024-00747-1
- Webster, A. D., Spencer, J. E., Dover, C., and Atkinson, C. J. (2006). "The influence of sprays of gibberellic acid (GA₃) and aminoethoxyvinylglycine (AVG) on fruit abscission, fruit ripening and quality of two sweet cherry cultivars," *Acta Horticulturae* 727, 467-472. DOI: 10.17660/ActaHortic.2006.727.57
- White, P. J., and Broadley, C. L. (2003). "Calcium in plants," *Ann. Bot.* 92(4), 487-511. DOI: 10.1093/aob/mcgl64
- Wood, R. M., de Freitas, S. T., Argenta, L. C., and Neuwald, D. A. (2024). "The influence of pre-harvest calcium application on the concentration and distribution of ascorbic acid and mineral content in apple cultivars at harvest and during storage," *Postharvest Biology and Technology* 214, article 112979. DOI: 10.1016/j.postharvbio.2024.112979
- Wu, Y., Dong, X., Zhu, Y., and Chen, H. (2025). "Influence of pre-harvest sprays of calcium chloride on fruit storage quality and softening of 'fengtangli' (*Prunus*

salicina L.) plum,” *CyTA – Journal of Food* 23(1), article 2446834. DOI: 10.1080/19476337.2025.2446834

Xu, T., Niu, J., and Jiang, Z. (2022). “Sensing mechanisms: Calcium signaling mediated abiotic stress in plants,” *Frontiers in Plant Science* 13, article 925863. DOI: 10.3389/fpls.2022.925863

Zahid, A., Razzaq, A. A., Munawar, M., Ramzan, M., Almutairi, B. O., and Almutairi, M. H. (2023). “Foliar spray of salicylic acid and ascorbic acid ameliorates the biochemical compounds in hybrid chillies,” *Journal of King Saud University - Science* 35(5), article 102660. DOI: 10.1016/j.jksus.2023.102660

Zhang, L., Wang, P., and Johnson, K. (2023). “Optimizing gibberellin application timing to enhance fruit size and firmness in ‘Bing’ sweet cherry,” *HortScience* 58(2), 215-222. DOI: 10.21273/HORTSCI16988-22

Article submitted: July 26, 2025; Peer review completed: October 5, 2025; Revised version received and accepted: October 7, 2025; Published: October 21, 2025.
DOI: 10.15376/biores.20.4.10504-10520